



---

## Research Article

# Microbiological Evaluation of Soil and Wastewater from Major Abattoir Sites in Umuahia, Abia State

Ekundayo, C. I.<sup>1</sup>, Okafor, A. T.<sup>1</sup>, and \*Enya, E.<sup>2</sup>

<sup>1</sup>Department of Environmental Management and Toxicology, College of Natural Resources and Environmental Management, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria

<sup>2</sup>Department of Microbiology, College of Natural Sciences Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria

\*Corresponding Author's email: [emmy\\_anya12@yahoo.com](mailto:emmy_anya12@yahoo.com); Phone: +2347068119871

---

## ABSTRACT

Abattoirs with their associated activities produces enormous amount of wastewater that can alter the microbiota of the immediate environment. This study evaluated the microbial profile of wastewater and soil within two major Abattoirs, in Umuahia, Abia state, Nigeria. Wastewater and soil samples within the immediate environment (at different depths) were collected aseptically from different location (Ubakala and Ndioru). The samples were inoculated into culture media for isolation of microorganisms and identified accordingly using standard techniques. The total heterotrophic bacterial count ranged from  $8.7 \times 10^6 - 1.58 \times 10^7$  cfu/g and  $8.1 \times 10^6 - 1.33 \times 10^7$  cfu/g for the on-site and offsite soil samples with the Ubakala samples significantly recording higher counts than Ndioru. The wastewater samples from both locations had counts in the range of  $9.6 \times 10^6 - 1.36 \times 10^7$  cfu/mL while coliform counts ranged between  $3.7 \times 10^4 - 5.3 \times 10^4$  cfu/mL. Diverse microorganisms were isolated as *Bacillus subtilis* *Lysinibacillus* species, *Pseudomonas aeruginosa*, CONs, *Micrococcus* species, *Escherichia. coli*, *Acetobacter* species and *Enterobacter* species while the fungal isolates included *Mucor* sp., *Rhizopus* sp. and *Penicillium* sp., *Curvularia* species, *Cladosporium* species, *Aspergillus fumigatus*. The results as obtained in this study revealed that bacterial and fungal diversity was largely higher in onsite samples compared to offsite samples. This is a clear indication of the direct influence of abattoir effluent on soil microbial communities. The contamination level especially with pathogenic species as seen from this study shows that the study sites could be breeding grounds conducive for silent transmission of pathogen.

**Keywords:** Abattoir; *Escherichia coli*; Pathogen; Wastewater

**Citation:** Ekundayo, C.I., Okafor, A.T., & Enya, E. (2026). Microbiological Evaluation of Soil and Wastewater from Major Abattoir Sites in Umuahia, Abia State. *Sahel Journal of Life Sciences FUDMA*, 4(2): 76-83. DOI: <https://doi.org/10.33003/sajols-2026-0402-10>

---

## INTRODUCTION

Abattoir wastewater usually has a complex composition that could be potentially detrimental to the environment and endanger public health (Emeh *et al.*, 2020). In many developing nations including Nigeria, abattoir business is an important component of the livestock industry, as it provides constant meat supply to people and employment opportunities for a good number of the population (Nafaranda *et al.*,

2012). However, this industry is ravaged with poor facilities for the treatment of abattoir effluents (Ogbonnaya, 2011). Thus, large amounts of solid waste and effluent such as rumen contents, blood and wastewater are generated from abattoirs and they pollute surface and ground water with pathogens and undesirable chemical compounds (Olaiya *et al.*, 2016).

Different types of animals are usually slaughtered in the abattoir, with their blood, part of the dung and abdominal content washed on cemented pavements. The animal blood and most other wastes are most often released untreated into the flowing stream while the consumable parts of the slaughtered animal are washed directly into the flowing water (Omole and Longe, 2008). The release of this wastewater into water bodies is of public health concern because the water bodies into which these effluents are discharged is used by the locals for innumerable purposes including being a source of potable water, domestic water and for irrigation of farm lands (Emeh *et al.*, 2020).

Abattoirs by their nature serve as breeding grounds for pathogens and contaminants, polluting the environment and threatening public health (Anele *et al.*, 2023). This is so because these wastes are generated from unscreened animals that are slaughtered in the abattoirs which are hosts to microorganisms including bacteria (Constance *et al.*, 2019). Microorganisms present in abattoir effluents may include *Cryptosporidium parvum*, *Campylobacter* spp., *Yersinia enterocolitica*, hepatitis E virus, *Salmonella* spp., rotaviruses, *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Giardia lamblia* which are known pathogens (Ubwa *et al.*, 2013). The discharge of effluents from abattoirs into a water body introduces some potentially pathogenic microorganisms into the water body which is used either for domestic, recreation or irrigation purposes thus constituting environmental and public health challenge. Several studies including those of Ovuru *et al.* (2024) as well as Fasanmi *et al.* (2017) have revealed that abattoirs in developing countries have an unhygienic environment and detected the presence of pathogens that are known causes of diarrheal diseases (Adeyemo, 2002; Nwanta *et al.*, 2010).

There are increased chances of portions of these wastewaters been washed directly to the soil within the vicinity which may affect microbial species diversity and buildup of pathogens in the food chain. This study hence, evaluated the microbiological profile of wastewater effluents generated from the abattoir as well as the surrounding soil.

## **MATERIALS AND METHODS**

### **Sample collection**

Wastewater and soil samples within the immediate environment (at different depths) were collected from abattoirs in two different locations (Ubakala and Ndioru) of Abia State into sterile bottles. The samples were immediately transported in an ice box to the laboratory for analyses within 6 hours of collection.

### **Media Preparation**

The culture media (Nutrient agar, MacConkey agar and Sabouraud dextrose agar) used for this study were prepared by weighing out specific grams of the agar powder and dissolving in equivalent volume of distilled water following the directives stipulated by the manufacturers of the medium. They were sterilized by autoclaving at 121°C for 15minutes, allowed to cool before pouring into plates.

### **Isolation of Microorganisms from the samples**

The serial dilution technique was employed in the inoculation of the soil/water samples. Each of the samples was diluted in the 10-fold serial dilution technique described by Gurung *et al.* (2009). Six test tubes were set up in a rack and filled with 9ml of sterile distilled water labelled stock,  $10^{-1}$ ,  $10^{-2}$ ,  $10^{-3}$ ,  $10^{-4}$  and  $10^{-5}$  respectively. One (1) g and 1ml each of the soil/water sample was added to the tube labelled stock and properly mixed together. An Aliquot (1ml) was transferred from this tube into the one labelled  $10^{-1}$  and carefully mixed. Similarly, 1ml was drawn up from the  $10^{-1}$  tube and transferred to the  $10^{-2}$  tube. This was continued up to the last tube labelled  $10^{-5}$  from where 1ml was discarded. Dilutions from  $10^{-3}$  and  $10^{-4}$  were inoculated onto freshly nutrient agar, MacConkey agar and Sabouraud Dextrose agar plates. The Spread plate method of inoculation as described by Prescott *et al.* (2007) was used where 0.1ml of the respective dilutions ( $10^{-3}$  and  $10^{-4}$ ) were plated on various agar plates and evenly spread over the entire plate using a flame sterilized glass rod. The inoculated plates were incubated at 35°C for 24hrs for bacteria and at room temperature ( $25\pm 2^{\circ}\text{C}$ ) for fungi.

### **Characterization and Identification of the Isolates**

#### **Gram staining**

The method of Green-Berge *et al.* (1992) was adopted in identification of bacteria Gram reaction (Gram positive and Gram negative). Bacteria smear was made on a clean grease-free slide, air dried and heat fixed. The slide was flooded with crystal violet for 1 minute, decanted and rinsed with water. Lugol's

iodine (mordant) was applied for 60 seconds and rinsed. Acetone was used in decolorizing and washed immediately then counter stained with neutral red for 1 minute. It was then rinsed with water, blotted carefully and air dried. Finally, the slides were observed under the microscope using oil immersion objectives (x100).

#### Biochemical Tests

Isolated organisms were identified by standard microbiology identification techniques including catalase test, citrate, hydrogen sulphide test, starch hydrolysis test, methyl-red test, voges-proskauer test, oxidase test as well as the urease and indole tests.

#### RESULTS

The results presented in Table 1 shows the microbial counts across different sample types (soil at two depths, wastewater, and clean water controls) for two abattoir locations (Ndoru and Ubakala). The total heterotrophic bacterial count (THBC) ranged from 8.7

$\times 10^6$  cfu/g –  $1.58 \times 10^7$  cfu/g and  $8.1 \times 10^6$  cfu/g –  $1.33 \times 10^7$  cfu/g for the on-site and offsite soil samples with the Ubakala samples significantly recording higher counts than Ndioru. Clean water controls had lower counts between  $3.8 \times 10^6$  cfu/mL –  $4.3 \times 10^6$  cfu/mL. Coliforms were detected in most onsite soils and wastewater samples, but absent in several offsite soils while the fungal load ranged from  $3.4 \times 10^4$  cfu/g –  $7.6 \times 10^4$  cfu/g at the Ndioru site and within  $2.8 \times 10^4$  cfu/g –  $6.8 \times 10^4$  cfu/g at Ubakala. The bacterial isolates recovered from the samples in this study were identified as *B. subtilis*, *Lysinibacillus* species, *Pseudomonas aeruginosa*, CONs, *Micrococcus* species, *E. coli*, *Acetobacter* species and *Enterobacter* species while the fungal isolates included *Mucor* sp., *Rhizopus* sp. and *Penicillium* sp., *Curvularia* species, *Cladosporium* species, *Aspergillus fumigatus*. The results as obtained in this study revealed that bacterial and fungal diversity was largely higher in onsite samples compared to offsite samples.

**Table 1: Total Heterotrophic bacterial and fungal counts (in cfu/g & cfu/ml) of the samples**

Samples	THBC	Coliform count	Fungal load
Ndoru@15cm offsite (Soil)	$8.1 \times 10^6$	NG	$3.4 \times 10^4$
Ndoru@10cm offsite (Soil)	$1.08 \times 10^7$	$1.8 \times 10^4$	$6.1 \times 10^4$
Ndoru@15cm onsite (Soil)	$8.7 \times 10^6$	$3.3 \times 10^4$	$3.8 \times 10^4$
Ndoru@10cm onsite (Soil)	$1.27 \times 10^7$	$4.1 \times 10^4$	$7.6 \times 10^4$
Wastewater Ndoru1	$1.08 \times 10^7$	$3.7 \times 10^4$	$3.8 \times 10^4$
Wastewater Ndoru2	$1.36 \times 10^7$	$5.3 \times 10^4$	$4.6 \times 10^4$
Clean water (Control)	$4.3 \times 10^6$	$1.8 \times 10^4$	$2.1 \times 10^4$
Ubakala@15cm offsite (Soil)	$1.19 \times 10^7$	NG	$2.8 \times 10^4$
Ubakala@10cm offsite (Soil)	$1.33 \times 10^7$	NG	$4.2 \times 10^4$
Ubakala@15cm onsite (Soil)	$1.03 \times 10^7$	NG	$3.8 \times 10^4$
Ubakala@10cm onsite (Soil)	$1.58 \times 10^7$	$4.6 \times 10^4$	$6.8 \times 10^4$
Wastewater Ubakala1	$1.12 \times 10^7$	$3.8 \times 10^4$	$3.8 \times 10^4$
Wastewater Ubakala2	$9.6 \times 10^6$	$4.6 \times 10^4$	$4.6 \times 10^4$
Clean water (Control)	$3.8 \times 10^6$	$2.6 \times 10^4$	NG

THBC: Total heterotrophic bacterial count; NG: No Growth

**Table 2: Macroscopic and Microscopic Description of Isolated Fungi**

Macroscopic Appearance	Microscopic features	Probable fungi
White fluffy colonies that turned grey with age	Long branching sporangiophores with non-septate hyphae bearing spherical spores at the tip	<i>Mucor</i> species
Bright-green colonies with white edges	Long ellipsoidal conidia with smooth walled conidiophores borne on septate hyphae	<i>Penicillium</i> species
Rapidly growing white cottony colonies on PDA plate	Upright sporangiophores borne on aseptate hyphae with numerous spiral spores.	<i>Rhizopus</i> species
Brown to blackish brown fluffy colonies with a black reverse pigmentation	Erect conidiophores borne on septate hyphae	<i>Curvularia</i> species
Colonies were dark olive and flat. The reverse side was black	Hyphae were septate, with erect Conidiophores bearing Conidia	<i>Cladosporium</i> species
Yellow or yellowish green colonies with distinct margin	Hyphae were septate, simple and thick-walled. Conidiophores bearing conidial heads containing conidia were seen	<i>Aspergillus fumigatus</i>

**Table 3: Frequency of Isolation of each isolate**

Bacterial Isolates	No. (%) Isolated	Fungal Isolates	No. (%) Isolated
<i>Micrococcus</i> species	6 (42.8)	<i>Cladosporium</i> species	4 (28.6)
<i>Bacillus subtilis</i>	4 (28.6)	<i>Aspergillus fumigatus</i>	7 (50.0)
Coagulase Negative Staphylococci (CONs)	9(64.2)	<i>Curvularia</i> species	6 (42.8)
<i>Pseudomonas aeruginosa</i>	4 (28.6)	<i>Penicillium</i> species	6 (42.8)
<i>Enterobacter</i> species	5 (35.7)	<i>Mucor</i> species	8 (57.1)
<i>Escherichia coli</i>	6 (42.8)	<i>Rhizopus</i> species	8 (35.7)
<i>Salmonella</i> species	6 (42.8)		
<i>Bacillus megaterium</i>	6 (42.8)		
<i>Lysinibacillus</i> species	5 (35.7)		
<i>Acetobacter</i> species	4 (28.6)		

**Table 4: Distribution of the Fungal isolates across the sample types**

Samples	<i>Rhizopus</i> species	<i>Penicillium</i> species	<i>Aspergillus fumigatus</i>	<i>Curvularia</i> species	<i>Mucor</i> species	<i>Cladosporium</i> species
A	-	+	+	-	+	-
B	+	+	+	-	+	+
C	+	-	+	-	-	-
D	+	+	-	+	+	-
E	-	-	+	+	+	-
F	-	-	-	-	+	-
G	-	-	-	+	-	-
H	-	+	+	-	-	-
I	+	-	+	-	-	+
J	-	+	-	-	+	-
K	+	+	-	+	-	+
L	-	-	-	+	+	+
M	-	-	+	+	+	-
N	-	-	-	-	-	-

A: Ndor@15cm offsite; B: Ndor@10cm offsite; C: Ndor@15cm onsite; D: Ndor@10cm onsite; E: Wastewater Ndor1; F: Wastewater Ndor2; G: Clean water Control Ndor; H: Ubakala@15cm offsite; I: Ubakala@10cm offsite; J: Ubakala@15cm onsite; K: Ubakala@10cm onsite; L: Wastewater Ubakala1; M: Wastewater Ubakala2; N: Clean water control Ubakala.

Table 5: Distribution of the Bacterial isolates across the sample types

Samples	<i>B. subtilis</i>	<i>Lysinibacillus</i> species	<i>Pseudomonas aeruginosa</i>	CONs	<i>Micrococcus</i> species	<i>E. coli</i>	<i>Acetobacter</i> species	<i>Enterobacter</i> species	<i>Salmonella</i> species
A	+	+	+	-	+	-	-	-	-
B	+	-	-	+	+	-	-	-	+
C	-	-	+	-	-	+	-	+	-
D	-	+	-	+	+	+	-	-	+
E	-	-	-	+	-	-	+	+	+
F	-	-	-	-	-	+	+	-	+
G	-	-	-	+	+	-	-	+	-
H	+	+	-	-	-	-	-	-	-
I	-	+	+	+	-	-	+	-	-
J	+	-	+	+	-	-	-	-	-
K	-	+	-	+	+	+	-	-	+
L	-	-	-	+	-	+	+	-	-
M	-	-	-	-	+	+	-	+	+
N	-	-	-	+	-	-	-	+	-

A: Ndoru@15cm offsite; B: Ndoru@10cm offsite; C: Ndoru@15cm onsite; D: Ndoru@10cm onsite; E: Wastewater Ndoru1; F: Wastewater Ndoru2; G: Clean water Control Ndoru; H: Ubakala@15cm offsite; I: Ubakala@10cm offsite; J: Ubakala@15cm onsite; K: Ubakala@10cm onsite; L: Wastewater Ubakala1; M: Wastewater Ubakala2; N: Clean water control Ubakala.

## DISCUSSION

The total heterotrophic bacterial count (THBC) ranged from  $8.7 \times 10^6$  cfu/g –  $1.58 \times 10^7$  cfu/g and  $8.1 \times 10^6$  cfu/g –  $1.33 \times 10^7$  cfu/g for the on-site and offsite soil samples with the Ubakala samples significantly recording higher counts than Ndioru. The wastewater samples from both locations had counts in the range of  $9.6 \times 10^6$  cfu/mL –  $1.36 \times 10^7$  cfu/mL while coliform counts ranged between  $3.7 \times 10^4$  cfu/mL –  $5.3 \times 10^4$  cfu/mL. Anele *et al.* (2023) reported a THBC of  $3.3 \times 10^5$  –  $5.0 \times 10^6$  cfu/g in a related study on environmental impact assessment of abattoirs in Rivers State, Nigeria. Equally, similar counts have been reported by (Nafarnda *et al.*, 2012; Ogbonnaya, 2011).

The total heterotrophic bacterial counts (THBC) in the soil samples at Ubakala recorded higher bacterial counts more than those of Ndioru probably due to the degree of movement and activity at the Ubakala site. Also, the abundance of microorganisms can be attributed to the abundance of nutrients in the wastewater, which are suitable for optimal microbial metabolism (Neboh *et al.*, 2013; Darbar and Saha, 2023). It can be seen that the value  $9.6 \times 10^6$  cfu/mL –  $1.36 \times 10^7$  cfu/mL was high, indicating that the wastewater ought to be treated before onward discharge into the environment. The counts as obtained in this study is in contrast to the WHO allowable limit ( $1 \times 10^2$  cfu/mL) reported by Dankaka *et al.* (2018). Rabah *et al.* (2008) reported a mean ( $6.4 \times 10^7$  cfu/mL) for the total viable count of bacteria isolated in Sokoto abattoir effluent, which is consistent with the result of our study.

The bacterial isolates recovered from the samples in this study were identified as *B. subtilis*, *Lysinibacillus* species, *Pseudomonas aeruginosa*, CONs, *Micrococcus* species, *E. coli*, *Acetobacter* species and *Enterobacter* species while the fungal isolates included *Mucor* sp., *Rhizopus* sp. and *Penicillium* sp., *Curvularia* species, *Cladosporium* species, *Aspergillus fumigatus*. Anele *et al.* (2023) also identified similar bacterial isolates in wastewater samples in abattoirs within Port Harcourt. The diversity of microorganisms as obtained in this study is in line with the results of Obire and Ariyo (2021), who identified *Salmonella* sp. (13.16%), *Bacillus* sp. (20.16%), *Pseudomonas* sp. (16.04%), *Escherichia coli* (12.34%), *Micrococcus* sp. (9.46%) and *Staphylococcus* sp. (22%), from Niger Delta abattoir soils.

The high frequency of CONs (64.2%) is particularly noteworthy. *Staphylococcus aureus* was also the most common isolate (28.5%) from Akure abattoirs,

according to Adegunloye (2013), demonstrating the prevalence of staphylococci in these kinds of environment. The presence of *Enterobacter* species (35.7%) and *Pseudomonas aeruginosa* (28.6%) is of significance, as both are capable of causing a range of infections. *Salmonella* species were found in 42.8% of samples from several sites in this study. This is a crucial finding with major consequences for environmental and public health. Ijah *et al.* (2022) also reported *Salmonella* as prevalent pathogens in Nigerian abattoir environments, with counts ranging from  $1.8 \times 10^3$  to  $2.1 \times 10^4$  cfu/mL in Minna abattoir wastewater. It was also found that at both onsite soils and wastewater, 42.8% of samples had *Escherichia coli*. This aligns with findings from Oladipo *et al.* (2022), who identified *E. coli* as a predominant organism in abattoir wastewater and contaminated soils. Equally, Akindele *et al.* (2015) reported *E. coli* as one of the key bacterial isolates from Ijebu-Igbo abattoir. The presence of these bacteria especially *Salmonella* species and *Escherichia coli* portends a grave public health concern as these microorganisms are linked with water borne diseases as they can persist in the environment and could contaminate nearby water sources through run-offs (Olaiya *et al.* 2016).

There were six different fungus genera identified in this study with *Aspergillus fumigatus* having the highest dominance (50%). The presence of this organism is significant as it is a known opportunistic pathogen that can cause aspergillosis in immunocompromised persons. Similar fungal profiles were found in Niger Delta abattoir soils, according to the findings of Obire and Ariyo (2021), with *Penicillium* (43.3%) being the most common isolate, followed by *Aspergillus* sp. (26.8%), *Fusarium* sp. (11.6%), and *Mucor* (9.2%). Related studies in different regions of Nigeria have recorded similar microbial profile especially those of Ogbomida *et al.*, (2016), Ogunnusi and Dahunsi (2014), Shukri *et al.* (2017) Njoku *et al.* (2018) and Awari *et al.* (2020) who also identified bacterial pathogens in effluent samples from abattoirs.

The results as obtained in this study revealed that bacterial and fungal diversity was largely higher in onsite samples compared to offsite samples. This is a clear indication of the direct influence of abattoir effluent on soil microbial communities and is consistent with findings from Ekpengkho *et al.* (2025), who reported increased bacterial and fungal counts in abattoir-impacted soils compared to control sites. Meanwhile, clean water controls showed limited microbial diversity, with only Coagulase negative

Staphylococcus (CONs) and *Micrococcus* detected at Ndor, and CONs alone at Ubakala affirming that the high microbial loads observed at abattoir sites are directly attributable to contamination from slaughter activities.

## CONCLUSION

The microbiological evaluation of soil and wastewater from Ndor and Ubakala abattoir sites reveals heavy contamination with diverse bacterial and fungal populations, including multiple pathogenic species. The contamination level especially with pathogenic species as seen from this study shows that the study sites could be breeding grounds conducive for silent pathogen transmission. It is thus suggested improved sanitation infrastructures, and enhanced surveillance around abattoir sites be enforced to preserve the environment and protect the public.

## REFERENCES

Adegunloye, D.V. (2013). Microbial composition of the abattoir environment and its health implications on the quality of fresh cow meat sold in Akure, Ondo State, Nigeria. *WIT Transactions on Ecology and the Environment*, 170, 57-65

Adesemoye, A.O. and Adedire, C. O. (2005). Use of cereals as basal medium for the formulation of alternative culture media for fungi. *World Journal of Microbiology and Biotechnology*, 21, 329-336.

Adeyemo, O. K., Ayodeji, I. O. and Aika-Raji, C. O. (2002). The water quality and sanitary conditions in a major Abattoir (Bodija) in Ibadan, Nigeria. *African Journal of Biomedical Research*, 5:1-2.

Akindele, S.T., Abimbola, W.A., Oluwabiyi, B.A., Jokotagba, O.A., Agbolade, O. M. & Sam-Wobo, S. O. (2015). Microbiological assessment of waste-water and soil in Ijebu-Igbo abattoir, Southwest, Nigeria. *International Journal of Advancements in Research & Technology*, 4(10)

Anele, B.C., Okerentuagha, P.O., Stanley, H.O. and Immanuel, O.M. (2023). Environmental impact assessment of abattoirs in Rivers State Nigeria. *World Journal of Advanced Research and Reviews*, 19(2), 1014-1023.

Anele, B.C., Okerentugba, P.O., Stanley, H.O., Immanuel, O.M., I.M., Ikeh, I.M., Ukanwa, C.C. and Okonko, I.O. (2023). Environmental impact assessment of abattoirs in Rivers State, Nigeria. *World Journal of Advanced Research and Reviews*, 19(02), 1014–1023

Awari, V.G., Owhouka, A. & Adanella, L.P. (2020). Assessment of mycological abattoir waste water

contaminated soil in Port Harcourt. *Journal of Advances in Microbiology Research*, 1(2), 21-28.

Constance, O. Egesi, C.O., Victor, E. Anthony, C.E. and Charles, E.O. (2019). Compliance Monitoring of Microbiological and Physicochemical Parameters of Abattoirs' Effluents Discharged into Water Bodies in Owerri, Nigeria. *Microbiology Research Journal International*. 28(6): 1-16.

Dankaka, S.M., Farouq, A.A., Bagega, A.I. & Abubakar, U. (2018). Microbiological and physicochemical analysis of old Sokoto abattoir wastewater. *Bioremediation Science and Technology Research Journal*, 6(2), 9-13.

Darbar, S. & Saha, S. (2023). Novel multi-herbal formulation (SS-IM-21) and its sanative effect in CCl<sub>4</sub> induced experimental animals. *International Journal of Pharmaceutical and Phytopharmacological Research*, 13(2), 30-36.

Deborah, J. M., Dammo, M. N., Watila, S. and Sangodoyin, A. Y. (2017). Impacts of Effluents on River Nggada water quality in Maiduguri, Nigeria. *Advancement in Scientific and Engineering Research*. 3(1):1-9.

Ekpenkhio, E., Ugwa, I. K. & Adebajo, O. D. (2025). Evaluating the impact of abattoir wastes on soil properties in Benin City, Edo State, Nigeria. *Ife Social Sciences Review*, 33(1), 173-186

Emeh, A.A., Anyanwu, G.O., Ibe, I.J., Emeh, T.C., Odaghara, C.J. and Nwaehiri, U.L. (2020). Assessment of the impact of EGBU abattoir effluent on the microbiological properties of Otamiri River. *Int J Sci Res Publ.*, 10:5.

Fasanmi, O. G., Ayodeji, I. O., Oloso, N. O. and Fasina, F. O. (2017). Retrospective studies of abattoir zoonoses in Nigeria: public health implications. *CAB Reviews*, 12(058), 1–14.

Green-Berge, L.S.H, Sahilah, A.M., Son, R. and Tosiah, S. (1992). Isolation and screening of actinomycetes from Malaysian soil for their enzymatic and antimicrobial activities. *Journal of Tropical Agriculture and Food Sciences*, 35: 159-164.

Gurung, T.D, Sherp, C., Agrawal, V.P. and Lekhak, B. (2009). Isolation and characterization of antibacterial actinomycetes from soil samples of Kalapatthar Mount Everest Region. *Nepal Journal of Science and Technology*, 10: 173-182.

Ijah, U. J. J., Musa, I. O., Amao, F., Maude, A. M. & Victor-Ekwebelem, M. O. (2022). Wastewater quality and unhygienic practices in Minna abattoir, North Central Nigeria. *Science World Journal*, 17(2)

Kazzah, R. C., Kutman, N. A., Michael, A. M., Obaje, B. K., Fodeke, H., Feka, D. P., Tanko, S. F. & Bammai, M. Y. (2020). Elemental and bacterial profile of soils

- contaminated by effluent originating from Zangon-Shanu Abattoir, Zaria: A preliminary investigation. *International Journal of Research and Innovation in Applied Science*, 7(11), 01-05
- Nafarnda, W. D., Ajayi, I. E., Shawulu, J. C., Kawe, M. S., Omeiza, G. K., Sani, N. A., Tenuche, O. Z., Dantong, D. D., and Tags, S. Z. (2012). Bacteriological Quality of Abattoir Effluents Discharged into Water Bodies in Abuja, Nigeria. *Veterinary Science*. Volume 2012, Article ID 515689, 5 pages.
- Ndakaru, O.E. & Ohwo, O. (2023). Seasonality assessment of abattoir waste impact on water quality of Anwai River in Nigeria. *Biodiversity Journal*, 14(3), 405-413.
- Njoku-Tony, R. F., Ogbuagu, D. H., Ihejirika, C. E., Nwoko, C. O., Amaku, G. E., Azoro, V. A., Ukaegbu, K., Ezikeudu, E. C., and Edafienene, E. O. (2018). Impact of Abattoir Waste on the Water Quality of Amilimocha River Asaba, Delta State. *International Journal of Energy and Environmental Research*. 6(1):25-35.
- Nwanta, J. A., Onunkwo, J. and Ezenduka, E. (2010). Analysis of Nsukka metropolitan abattoir solid waste and its bacterial contents in south eastern Nigeria: public health implication. *Archives of Environmental & Occupational Health*, 65(1), 21–26.
- Obire, O. & Ariyo, A. B. (2021). An evaluation of microbiological and physicochemical characteristics of abattoir soils in the Niger Delta. *International Journal of Research and Innovation in Applied Science*, 6(9), 126-133
- Ogbomida, E. T., Kubeyinje, B. & Ezemonye, L. I. (2016). Evaluation of bacterial profile and biodegradation potential of abattoir wastewater. *African Journal of Environmental Science and Technology*, 10(2), 50-57.
- Ogbonnaya, C. (2011). Analysis of groundwater pollution from abattoir waste in Minna, Nigeria. *Research Journal of Dairy Science*, 2(4): 74-77
- Ogunnusi, T.A. & Dahunsi, O.V. (2014). Isolation and identification of microorganisms from abattoir effluents from Oyo, Oyo State, Nigeria. *Asian Journal Applied Science*, 2(2), 218-222.
- Ojekunle, O.Z. and Lateef, S.T. (2017). Environmental Impact of Abattoir Waste Discharge on the Quality of Surface Water and Ground Water in Abeokuta. *Journal of Environmental and Analytical Toxicology*; 7: 509.
- Oladipo, O. O., Ilesanmi, T. M. & Olaleye, A. C. (2022). Environmental impact of abattoir waste discharge on the quality of water body and soil in Ado-Ekiti, Ekiti State. *Asian Journal of Biotechnology and Bioresource Technology*, 8(2), 1-6
- Olaiya, S., Mahmud, H., Eboreime, L. and Afolabi, O. C. (2016). Physico-Chemical and Microbial Analysis of the Effects of Abattoirs Operation in Estako-West and Central, Edo-State, Nigeria. *Global Journal of Science Frontier Research Environment & Earth Science*, 16(3) 1-11
- Osibanjo, O. and Adie, G.U. (2007). Impact of effluent from Bodija abattoir on the physicochemical parameters of Oshunkaye stream in Ibadan City, Nigeria. *Afr J Biotechnol.*, 6(15):1806-11.
- Ovuru, K. F., Izah, S. C., Ogidi, O. I., Imarhiagbe, O., & Ogwu, M. C. (2024). Slaughterhouse facilities in developing nations: sanitation and hygiene practices, microbial contaminants and sustainable management system. *Food Science and Biotechnology*, 33(3), 519–537.
- Prescott, L.M., Harley, J.P. and Klein, D.A. (2007). *Microbiology*, (7th Edition). London: McGraw Hill Publishers. P. 658
- Rabah, A., Ijah, U. J. J., Manga, S. B., & Ibrahim, M. L. (2008). Assessment of physicochemical qualities of abattoir wastewater in Sokoto, Nigeria. *Nigerian Journal of Basic and Applied Science*, 16(2), 149-154.
- Shukri, A. A., Kyambadde, J. and Hawumba, J. F. (2017). The Impact of Kalerwe Abattoir Wastewater Effluent on the Water Quality of the Nsooba Channel. *Agricultural Research and Technology*, 6:1
- Sumayya, B. U., Usman, B. U., Aisha, U., Shahida, A., Mohammad, A., Yakubu, M. S. and Zainab, M. (2013). Determination of Physiochemical Qualities of Abattoir Effluent on Soil and Water in Gandu, Sokoto State. *IOSR Journal of Environmental Science. Toxicology and Food*, 4(4): 47-50
- Tekenah, W. E., Agi, P. I. and Babatunde, B. B. (2014). Analysis of surface water pollution from abattoirs and the interrelationship between physico-chemical properties (A case study of the New Calabar River). *IOSR Journal of Environmental Science. Toxicology and Food Technology*, 8(5):10-18.
- Ubwa, S. T., Ato, G. H., Offem, J. O., Abah, J. and Asemave, K. (2013). An assessment of surface water pollution status around Gboko abattoir. *African Journal of Pure and Applied Chemistry*. 7(3): 131-138.
- Williams, J.O. and Dimbu, P.C. (2015). Effect of Abattoir Waste Water on Soil Microbial Communities. *Sch. Acad. J. Biosci.*, 3(5):452-455.